

TD 3: WEAK FORMULATION OF ELLIPTIC EQUATIONS

EXERCISE 1 (Ellipticity). For each of the following linear differential operator L , give the symbol, the principal symbol of L , and discuss the ellipticity and uniform ellipticity.

1. $Lu(x) = -\sum_{i,j=1}^d a_{ij}(x) \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i=1}^d b_i(x) \frac{\partial u}{\partial x_i} + c(x)u$, $x \in \Omega \subset \mathbb{R}^d$,
2. $Lf(x, v) = v \cdot \nabla_x f + F(x) \cdot \nabla_v f$, $x, v \in \mathbb{R}^d$, $F: \mathbb{R}^d \rightarrow \mathbb{R}^d$,
3. $Lu(t, x) = \partial_t u - \Delta u$, $t > 0$, $x \in \mathbb{R}^d$,
4. $Lu(t, x) = \partial_t u - i\Delta u$, $t > 0$, $x \in \mathbb{R}^d$.

EXERCISE 2 (Faber-Krahn inequality). Let Ω be an open bounded subset of \mathbb{R}^d with $d \geq 3$ and $V \in L^\infty(\Omega)$ such that $V \geq 0$. We consider the problem

$$(1) \quad \begin{cases} -\Delta u = Vu & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

1. Give the definition of a weak solution to (1).
2. Can you apply the Lax-Milgram theorem here?
3. Let $r > \frac{d}{2}$. Show that there is a constant $c_d > 0$ depending on d only such that, if (1) has a non-trivial weak solution, then

$$|\Omega|^{\frac{2}{d}-\frac{1}{r}} \|V\|_{L^r(\Omega)} \geq c_d.$$

Hint: Use the following Sobolev inequality

$$\|u\|_{L^{2^*}(\Omega)} \leq M_d \|\nabla u\|_{L^2(\Omega)}, \quad \frac{1}{2^*} = \frac{1}{2} - \frac{1}{d},$$

which holds for all $u \in H_0^1(\Omega)$, where M_d depends on d only.

4. What do you obtain in the particular case $V = \lambda = \text{cst}$?

EXERCISE 3 (Dirichlet problem). Let Ω be an open bounded subset of \mathbb{R}^d , $f \in L^2(\Omega)$ and $F \in L^2(\Omega)^d$. Show that the following elliptic problem with Dirichlet boundary condition

$$\begin{cases} -\Delta u = f - \text{div } F & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

has a unique weak solution $u \in H_0^1(\Omega)$.

EXERCISE 4 (Neumann problem). Let Ω be an open bounded subset of \mathbb{R}^d with smooth boundary, the exterior unit normal being denoted by n , and $f \in L^2(\Omega)$. Show that, for all $\mu > 0$, the elliptic problem with Neumann boundary condition

$$(2) \quad \begin{cases} -\Delta u + \mu u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega, \end{cases}$$

has a unique weak solution $u \in H^1(\Omega)$. In the case $\mu = 0$, give a necessary condition on $\int_\Omega f$ to the existence of a weak solution to (2).

EXERCISE 5 (Fourier condition). Let $\Omega \subset \mathbb{R}^d$ be an open bounded set with smooth boundary, $f \in L^2(\Omega)$, $g \in L^2(\partial\Omega)$ and $\lambda > 0$. We consider the following elliptic problem with Fourier boundary condition

$$(3) \quad \begin{cases} -\Delta u = f & \text{in } \Omega, \\ \lambda u + \frac{\partial u}{\partial n} = g & \text{on } \partial\Omega. \end{cases}$$

1. Give the variational formulation of the problem (3).
2. Prove that there exists a positive constant $C_\Omega > 0$ only depending on Ω such that for all $u \in H^1(\Omega)$,

$$\|u\|_{L^2(\Omega)}^2 \leq C_\Omega (\|\nabla u\|_{L^2(\Omega)}^2 + \lambda \|\gamma_0 u\|_{L^2(\partial\Omega)}^2),$$

where γ_0 denotes the trace operator $\gamma_0 : H^1(\Omega) \rightarrow L^2(\partial\Omega)$.

3. Prove that (3) has a unique weak solution.
4. * Is this weak solution a strong solution ?

EXERCISE 6 (The method of continuity).

1. Solve the equation $u - \Delta u = f$ on \mathbb{T}^d and show that it defines a map $L^2(\mathbb{T}^d) \rightarrow H^2(\mathbb{T}^d)$.
2. Let X, Y be some Banach spaces. Let $(T_t)_{t \in [0,1]}$ be a *continuous* path of continuous linear operators from X to Y satisfying

$$(4) \quad \exists C \geq 0, \forall u \in X, \forall t \in [0, 1], \quad \|u\|_X \leq C \|T_t u\|_Y.$$

Prove that T_0 is onto if and only if T_1 is onto as well.

3. Let $A \in C^1(\mathbb{T}^d, M_d(\mathbb{R}))$. We assume that the following ellipticity condition holds

$$\exists \alpha \in (0, 1), \forall x \in \mathbb{T}^d, \forall \xi \in \mathbb{R}^d, \quad A(x)\xi \cdot \xi \geq \alpha |\xi|^2.$$

We define the path $(T_t)_{t \in [0,1]}$ of operators $H^2(\mathbb{T}^d) \rightarrow L^2(\mathbb{T}^d)$ by the formula

$$T_t u = u - \operatorname{div}(A^{(t)}(x)\nabla u), \quad A^{(t)} = tA + (1-t)I_d.$$

- (a) Show that $t \mapsto T_t$ is continuous.
- (b) Check that (4) is satisfied.
- (c) Conclude.

EXERCISE 7 (Resolution by minimization). Let $\Omega \subset \mathbb{R}^3$ be open, bounded with smooth boundary. The purpose is to prove that the following elliptic problem has a non-trivial weak solution

$$\begin{cases} -\Delta u = u^3 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

1. Prove that there exists a solution to the following minimization problem

$$(5) \quad \inf \{ \|\nabla v\|_{L^2(\Omega)} : v \in H_0^1(\Omega), \|v\|_{L^4(\Omega)} = 1 \}.$$

Recall: Since $d = 3$ here, the continuous embedding $H_0^1(\Omega) \hookrightarrow L^q(\Omega)$ holds for all $1 \leq q \leq 6$, and is moreover compact when $1 \leq q < 6$.

2. Prove that if the function $v \in H_0^1(\Omega)$ solves (5), there exists a positive constant $\lambda > 0$ such that $-\Delta v = \lambda v^3$ weakly in Ω .
3. Conclude.