

Motivic knot theory

Clémentine Lemarié--Rieusset (Université de Bourgogne)

October 26, 2022

Algebraic geometry in a nutshell

Let F be a (perfect) field. Geometrical objects of interest are subsets of F^n which are zeroes of polynomials or complements of such subsets, for instance:

- F^n (no polynomial);
- the unit circle $\{(x, y) \in F^2, x^2 + y^2 - 1 = 0\}$ (1 p.);
- the diagonal line $\{(x, y) \in F^2, x - y = 0\}$ (1 p.);
- their intersection $\{(x, y) \in F^2, x^2 + y^2 - 1 = 0, x - y = 0\}$ (2 p.);
- the origin $\{0\} \subset F^2$: $\{(x, y) \in F^2, x = 0, y = 0\}$ (2 p.);
- $F^2 \setminus \{0\}$...

In practice, we replace these with schemes, for instance F^n is replaced with the affine n -space \mathbb{A}_F^n and $F^n \setminus \{0\}$ is replaced with the scheme $\mathbb{A}_F^n \setminus \{0\}$.

Knot theory in a nutshell

Topological objects of interest are knots and links.

- A **knot** is a (closed) topological subspace of the 3-sphere \mathbb{S}^3 which is homeomorphic to the circle \mathbb{S}^1 .
- An **oriented knot** is a knot with a “continuous” local trivialization of its tangent bundle, or equivalently of its normal bundle (the ambient space being oriented). There are two orientation classes.
- A **link** is a finite union of disjoint knots. A link is **oriented** if all its components (i.e. its knots) are oriented.
- The **linking number** of an (oriented) link with two components is the number of times one of the components turns around the other component.

Oriented knots and links in algebraic geometry

Recall that for all $n \geq 1$, \mathbb{S}^n has the same homotopy type as $\mathbb{R}^{n+1} \setminus \{0\}$.

Oriented knots and links in algebraic geometry

Recall that for all $n \geq 1$, \mathbb{S}^n has the same homotopy type as $\mathbb{R}^{n+1} \setminus \{0\}$.

Knot

A knot is a closed immersion $\varphi : \mathbb{A}_{\mathbb{F}}^2 \setminus \{0\} \rightarrow \mathbb{A}_{\mathbb{F}}^4 \setminus \{0\}$.

Oriented knots and links in algebraic geometry

Recall that for all $n \geq 1$, \mathbb{S}^n has the same homotopy type as $\mathbb{R}^{n+1} \setminus \{0\}$.

Knot

A knot is a closed immersion $\varphi : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$.

Link with two components

A link with two components is a couple of knots $\varphi_i : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$ with disjoint images Z_i (where $i \in \{1, 2\}$).

Oriented knots and links in algebraic geometry

Recall that for all $n \geq 1$, \mathbb{S}^n has the same homotopy type as $\mathbb{R}^{n+1} \setminus \{0\}$.

Knot

A knot is a closed immersion $\varphi : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$.

Link with two components

A link with two components is a couple of knots $\varphi_i : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$ with disjoint images Z_i (where $i \in \{1, 2\}$).

An orientation o_i of Z_i is a “trivialization” of the normal sheaf of Z_i in $\mathbb{A}_F^4 \setminus \{0\}$ (actually of its determinant (i.e. its maximal exterior power)).

Oriented knots and links in algebraic geometry

Recall that for all $n \geq 1$, \mathbb{S}^n has the same homotopy type as $\mathbb{R}^{n+1} \setminus \{0\}$.

Knot

A knot is a closed immersion $\varphi : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$.

Link with two components

A link with two components is a couple of knots $\varphi_i : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$ with disjoint images Z_i (where $i \in \{1, 2\}$).

An orientation o_i of Z_i is a “trivialization” of the normal sheaf of Z_i in $\mathbb{A}_F^4 \setminus \{0\}$ (actually of its determinant (i.e. its maximal exterior power)).

Intuition

Think of the normal sheaf of Z_i in $\mathbb{A}_F^4 \setminus \{0\}$ as a two-dimensional vector space and think of a trivialization of it as a basis of this vector space.

Orientation classes

Fact

The orientation classes are parametrized by the elements of $F^*/(F^*)^2$ (where $(F^*)^2 = \{a \in F^*, \exists b \in F^*, a = b^2\}$).

Orientation classes

Fact

The orientation classes are parametrized by the elements of $F^*/(F^*)^2$ (where $(F^*)^2 = \{a \in F^*, \exists b \in F^*, a = b^2\}$).

If $F = \mathbb{R}$ then $F^*/(F^*)^2$ has two elements.

Orientation classes

Fact

The orientation classes are parametrized by the elements of $F^*/(F^*)^2$ (where $(F^*)^2 = \{a \in F^*, \exists b \in F^*, a = b^2\}$).

If $F = \mathbb{R}$ then $F^*/(F^*)^2$ has two elements.

If $F = \mathbb{C}$ then $F^*/(F^*)^2$ has one element.

Orientation classes

Fact

The orientation classes are parametrized by the elements of $F^*/(F^*)^2$ (where $(F^*)^2 = \{a \in F^*, \exists b \in F^*, a = b^2\}$).

If $F = \mathbb{R}$ then $F^*/(F^*)^2$ has two elements.

If $F = \mathbb{C}$ then $F^*/(F^*)^2$ has one element.

If $F = \mathbb{Q}$ then $F^*/(F^*)^2$ has infinitely many elements (the classes of the integers without square factors).

The Hopf link

We fix coordinates x, y, z, t for \mathbb{A}_F^4 and u, v for \mathbb{A}_F^2 once and for all.

- The image of the Hopf link:

$$\{x = 0, y = 0\} \sqcup \{z = 0, t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

The Hopf link

We fix coordinates x, y, z, t for \mathbb{A}_F^4 and u, v for \mathbb{A}_F^2 once and for all.

- The image of the Hopf link:

$$\{x = 0, y = 0\} \sqcup \{z = 0, t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

- The parametrization of the Hopf link:

$$\varphi_1 : (x, y, z, t) \leftrightarrow (0, 0, u, v), \varphi_2 : (x, y, z, t) \leftrightarrow (u, v, 0, 0)$$

The Hopf link

We fix coordinates x, y, z, t for \mathbb{A}_F^4 and u, v for \mathbb{A}_F^2 once and for all.

- The image of the Hopf link:

$$\{x = 0, y = 0\} \sqcup \{z = 0, t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

- The parametrization of the Hopf link:

$$\varphi_1 : (x, y, z, t) \leftrightarrow (0, 0, u, v), \varphi_2 : (x, y, z, t) \leftrightarrow (u, v, 0, 0)$$

- The orientation of the Hopf link:

$$\sigma_1 : \bar{x}^* \wedge \bar{y}^* \mapsto 1, \sigma_2 : \bar{z}^* \wedge \bar{t}^* \mapsto 1$$

A variant of the Hopf link

- The image is the same as the Hopf link's image:

$$\{x = y, y = 0\} \sqcup \{z = 0, a \times t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\} \text{ with } a \in F^*$$

A variant of the Hopf link

- The image is the same as the Hopf link's image:

$$\{x = y, y = 0\} \sqcup \{z = 0, a \times t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\} \text{ with } a \in F^*$$

- The parametrization is the same:

$$\varphi_1 : (x, y, z, t) \leftrightarrow (0, 0, u, v), \varphi_2 : (x, y, z, t) \leftrightarrow (u, v, 0, 0)$$

A variant of the Hopf link

- The image is the same as the Hopf link's image:

$$\{x = y, y = 0\} \sqcup \{z = 0, a \times t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\} \text{ with } a \in F^*$$

- The parametrization is the same:

$$\varphi_1 : (x, y, z, t) \leftrightarrow (0, 0, u, v), \varphi_2 : (x, y, z, t) \leftrightarrow (u, v, 0, 0)$$

- The orientation is different:

$$\sigma_1 : \overline{x - y}^* \wedge \overline{y}^* \mapsto 1, \sigma_2 : \overline{z}^* \wedge \overline{at}^* \mapsto 1$$

Motivic homotopy theory

Overview

Motivic homotopy theory (a.k.a. \mathbb{A}^1 -homotopy theory) is a homotopy theory on smooth schemes of finite type over a “nice” base scheme (in our case the perfect field F).

The idea is to replace the unit interval $[0, 1]$ with the affine line \mathbb{A}_F^1 .

References on motivic homotopy theory

- The foundations were laid out in Morel and Voevodsky's article *\mathbb{A}^1 -homotopy theory of schemes* (1999)
- Its specificities when the base scheme is a perfect field were laid out in Morel's book *\mathbb{A}^1 -algebraic topology over a field* (2012)
- The nLab page *Motivic homotopy theory* is nicely done and has plenty of references

Motivic spheres

There are two analogues of the circle $[0, 1]/\{0, 1\}$ in motivic homotopy theory: $S^1 := \mathbb{A}_F^1/\{0, 1\}$ and the multiplicative group $\mathbb{G}_m := \mathbb{G}_{m,F}$.

Motivic spheres

For all $i, j \in \mathbb{Z}$, we denote by S^i the i -th smash-product of S^1 and we call the smash-product $S^i \wedge \mathbb{G}_m^{\wedge j}$ (in the stable homotopy category) a motivic sphere.

Note that the projective line $\mathbb{P}^1 := \mathbb{P}_F^1$ is equal to $S^1 \wedge \mathbb{G}_m$ in the stable homotopy category.

Intuition

Think of \mathbb{P}^1 as the set of lines in F^2 , i.e. $\{[x : y], (x, y) \in F^2 \setminus \{0\}\}$ with $[\lambda x : \lambda y] = [x : y]$ for all $\lambda \in F^*$.

Morel's Theorem

Objects of interest

The groups of morphisms $[S^i \wedge \mathbb{G}_m^{\wedge j}, S^k \wedge \mathbb{G}_m^{\wedge l}] = [S^{i-k}, \mathbb{G}_m^{\wedge(l-j)}]$ in the stable homotopy category.

Similarly to the fact that the stable homotopy group $\pi_i^s(S_0) = 0$ if $i < 0$, the group $[S^i, \mathbb{G}_m^{\wedge j}]$ is equal to 0 if $i < 0$ (with $j \in \mathbb{Z}$).

Morel's theorem

Morel gave a presentation by generators and relations of the graded ring with unit $\bigoplus_{n \in \mathbb{Z}} [S^0, \mathbb{G}_m^{\wedge n}]$ (where the product is given by the smash-product).

The generators are denoted $[a] \in [S^0, \mathbb{G}_m]$ for every $a \in F^*$ and $\eta \in [S^0, \mathbb{G}_m^{\wedge(-1)}] = [\mathbb{A}^2 \setminus \{0\}, \mathbb{P}^1]$ which sends (x, y) to $[x : y]$.

Milnor-Witt K -theory

Definition

The graded ring with unit $K_*^{\text{MW}}(F) := \bigoplus_{n \in \mathbb{Z}} [S^0, \mathbb{G}_m^{\wedge n}]$ is called the Milnor-Witt K -theory ring of F . We denote $K_n^{\text{MW}}(F) := [S^0, \mathbb{G}_m^{\wedge n}]$.

We denote $\langle a \rangle = \eta[a] + 1 \in K_0^{\text{MW}}(F)$ for every $a \in F^*$.

Fact

If $n \leq 0$ then every element of $K_n^{\text{MW}}(F)$ is a \mathbb{Z} -linear combination of $\langle a \rangle \eta^{-n}$ with $a \in F^*$.

The Rost-Schmid ring:

An analogue of the singular cohomology ring

To a smooth F -scheme Y , an integer $j \in \mathbb{Z}$ and an invertible \mathcal{O}_Y -module \mathcal{L} we associate the corresponding Rost-Schmid complex

$$\bigoplus_{i \geq 0} \bigoplus_{p \text{ point of codim } i \text{ in } Y} K_{j-i}^{\text{MW}}(\kappa(p)) \otimes \text{a twist which depends on } p \text{ and } \mathcal{L}.$$

The Rost-Schmid ring: An analogue of the singular cohomology ring

To a smooth F -scheme Y , an integer $j \in \mathbb{Z}$ and an invertible \mathcal{O}_Y -module \mathcal{L} we associate the corresponding Rost-Schmid complex

$$\bigoplus_{i \geq 0} \bigoplus_{p \text{ point of codim } i \text{ in } Y} K_{j-i}^{\text{MW}}(\kappa(p)) \otimes \text{a twist which depends on } p \text{ and } \mathcal{L}.$$

For every $i \in \mathbb{N}_0$, we denote the i -th cohomological group of this complex (called a Rost-Schmid group) by $H^i(Y, \underline{K}_j^{\text{MW}}\{\mathcal{L}\})$. We denote $H^i(Y, \underline{K}_j^{\text{MW}}) := H^i(Y, \underline{K}_j^{\text{MW}}\{\mathcal{O}_Y\})$.

The Rost-Schmid ring:

An analogue of the singular cohomology ring

To a smooth F -scheme Y , an integer $j \in \mathbb{Z}$ and an invertible \mathcal{O}_Y -module \mathcal{L} we associate the corresponding Rost-Schmid complex

$$\bigoplus_{i \geq 0} \bigoplus_{p \text{ point of codim } i \text{ in } Y} K_{j-i}^{\text{MW}}(\kappa(p)) \otimes \text{a twist which depends on } p \text{ and } \mathcal{L}.$$

For every $i \in \mathbb{N}_0$, we denote the i -th cohomological group of this complex (called a Rost-Schmid group) by $H^i(Y, \underline{K}_j^{\text{MW}}\{\mathcal{L}\})$. We denote $H^i(Y, \underline{K}_j^{\text{MW}}) := H^i(Y, \underline{K}_j^{\text{MW}}\{\mathcal{O}_Y\})$.

We have an intersection product

$$\cdot : H^i(Y, \underline{K}_j^{\text{MW}}) \times H^{i'}(Y, \underline{K}_{j'}^{\text{MW}}) \rightarrow H^{i+i'}(Y, \underline{K}_{j+j'}^{\text{MW}})$$

which makes $\bigoplus_{i \in \mathbb{N}_0, j \in \mathbb{Z}} H^i(Y, \underline{K}_j^{\text{MW}})$ into a graded $K_0^{\text{MW}}(F)$ -algebra.

Boundary maps

Definition

A boundary triple is a 5-tuple (Z, i, X, j, U) , or abusively a triple (Z, X, U) , with $i : Z \rightarrow X$ a closed immersion and $j : U \rightarrow X$ an open immersion such that the image of U by j is the complement in X of the image of Z by i , where Z, X, U are smooth F -schemes of pure dimensions. The boundary map associated to this boundary triple is the morphism

$$\partial : \mathcal{C}^\bullet(U, \underline{K}_*^{\text{MW}}) \rightarrow \mathcal{C}^{\bullet+1+d_Z-d_X}(Z, \underline{K}_{*+d_Z-d_X}^{\text{MW}}\{\nu_Z\})$$

induced by the differential d of the Rost-Schmid complex $\mathcal{C}(X, \underline{K}_*^{\text{MW}})$, i.e.:

$$\partial = i^* \circ d \circ j_*$$

The localization long exact sequence:

An analogue of the cohomology long exact sequ. of a pair

Theorem

Let (Z, i, X, j, U) be a boundary triple. The boundary map induces a morphism $\partial : H^{n+d_X-d_Z}(U, \underline{K}_{m+d_X-d_Z}^{\text{MW}}) \rightarrow H^{n+1}(Z, \underline{K}_m^{\text{MW}}\{\nu_Z\})$ and we have the following long exact sequence, called the localization long exact sequence:

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^n(Z, \underline{K}_m^{\text{MW}}\{\nu_Z\}) & \xrightarrow{i_*} & H^{n+d_X-d_Z}(X, \underline{K}_{m+d_X-d_Z}^{\text{MW}}) & \xrightarrow{j^*} & \dots \\ & & & & & & \\ & & \xrightarrow{j^*} & H^{n+d_X-d_Z}(U, \underline{K}_{m+d_X-d_Z}^{\text{MW}}) & \xrightarrow{\partial} & H^{n+1}(Z, \underline{K}_m^{\text{MW}}\{\nu_Z\}) & \longrightarrow \dots \end{array}$$

Punctured affine spaces are analogues of spheres

Let $n \geq 2$ be an integer, $i \in \mathbb{N}_0, j \in \mathbb{Z}$. The Rost-Schmid group $H^i(\mathbb{A}_F^n \setminus \{0\}, \underline{K}_j^{\text{MW}})$ is isomorphic to
$$\begin{cases} K_j^{\text{MW}}(F) & \text{if } i = 0 \\ K_{j-n}^{\text{MW}}(F) & \text{if } i = n - 1. \\ 0 & \text{otherwise} \end{cases}$$

This is similar to the fact in classical homotopy theory that $H^i(\mathbb{S}^{n-1})$ is isomorphic to
$$\begin{cases} \mathbb{Z} & \text{if } i = 0 \\ \mathbb{Z} & \text{if } i = n - 1. \\ 0 & \text{otherwise} \end{cases}$$

Note that $\mathbb{A}_F^n \setminus \{0\} = S^{n-1} \wedge \mathbb{G}_m^{\wedge n}$ in the stable homotopy category.

The linking number and its analogue

Let $L = K_1 \sqcup K_2$ be an oriented link (in knot theory) and \mathcal{L} be an oriented link with two components (in motivic knot theory), i.e. a couple of closed immersions $\varphi_i : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$ with disjoint images Z_i and orientation classes \overline{o}_i . We denote $Z := Z_1 \sqcup Z_2$.

The linking number and its analogue: step 1

Knot theory

The class S_i in $H^1(\mathbb{S}^3 \setminus L) \simeq H_2^{\text{BM}}(\mathbb{S}^3, L)$ of Seifert surfaces of the oriented knot K_i is the unique class that is sent by the boundary map to the (oriented) fundamental class of K_i in $H^0(K_i) \subset H^0(L)$.

The linking number and its analogue: step 1

Knot theory

The class S_i in $H^1(\mathbb{S}^3 \setminus L) \simeq H_2^{\text{BM}}(\mathbb{S}^3, L)$ of Seifert surfaces of the oriented knot K_i is the unique class that is sent by the boundary map to the (oriented) fundamental class of K_i in $H^0(K_i) \subset H^0(L)$.

Motivic knot theory

We define an analogue $[o_i] \in H^0(Z_i, \underline{K}_{-1}^{\text{MW}}\{\nu_{Z_i}\})$ of the oriented fundamental class of each oriented component of \mathcal{L} then we define the Seifert class \mathcal{S}_i as the unique class in $H^1(X \setminus Z, \underline{K}_1^{\text{MW}})$ that is sent by the boundary map to the oriented fundamental class $[o_i] \in H^0(Z, \underline{K}_{-1}^{\text{MW}}\{\nu_Z\})$.

The linking number and its analogue: step 2

Knot theory

The linking class of L is the image of the cup-product $S_1 \cup S_2 \in H^2(\mathbb{S}^3 \setminus L)$ by the boundary map $\partial : H^2(\mathbb{S}^3 \setminus L) \rightarrow H^3(\mathbb{S}^3, \mathbb{S}^3 \setminus L) \simeq H^1(L)$.

The linking number and its analogue: step 2

Knot theory

The linking class of L is the image of the cup-product $\mathcal{S}_1 \cup \mathcal{S}_2 \in H^2(\mathbb{S}^3 \setminus L)$ by the boundary map $\partial : H^2(\mathbb{S}^3 \setminus L) \rightarrow H^3(\mathbb{S}^3, \mathbb{S}^3 \setminus L) \simeq H^1(L)$.

Motivic knot theory

We define the quadratic linking class of \mathcal{L} as the image of the intersection product $\mathcal{S}_1 \cdot \mathcal{S}_2 \in H^2(X \setminus Z, \underline{K}_2^{\text{MW}})$ by the boundary map $\partial : H^2(X \setminus Z, \underline{K}_2^{\text{MW}}) \rightarrow H^1(Z, \underline{K}_0^{\text{MW}}\{\nu_Z\})$.

The linking number and its analogue: step 3

Knot theory

The linking number of $L = K_1 \sqcup K_2$ is the integer $n \in \mathbb{Z}$ such that the linking class in $H^1(L) = \mathbb{Z}[\omega_{K_1}] \oplus \mathbb{Z}[\omega_{K_2}]$ is equal to $(n[\omega_{K_1}], -n[\omega_{K_2}])$ (where ω_{K_i} is the volume form of the oriented knot K_i).

The linking number and its analogue: step 3

Knot theory

The linking number of $L = K_1 \sqcup K_2$ is the integer $n \in \mathbb{Z}$ such that the linking class in $H^1(L) = \mathbb{Z}[\omega_{K_1}] \oplus \mathbb{Z}[\omega_{K_2}]$ is equal to $(n[\omega_{K_1}], -n[\omega_{K_2}])$ (where ω_{K_i} is the volume form of the oriented knot K_i).

Motivic knot theory

We define the quadratic linking degree of \mathcal{L} as the image of the quadratic linking class of \mathcal{L} by the isomorphism

$$H^1(Z, \underline{K}_0^{\text{MW}} \{ \nu_Z \}) \rightarrow H^1(Z, \underline{K}_0^{\text{MW}}) \rightarrow H^1(\mathbb{A}_F^2 \setminus \{0\}, \underline{K}_0^{\text{MW}}) \oplus H^1(\mathbb{A}_F^2 \setminus \{0\}, \underline{K}_0^{\text{MW}}) \rightarrow K_{-2}^{\text{MW}}(F) \oplus K_{-2}^{\text{MW}}(F).$$

We fixed an isomorphism $H^1(\mathbb{A}_F^2 \setminus \{0\}, \underline{K}_0^{\text{MW}}) \rightarrow K_{-2}^{\text{MW}}(F)$ once and for all. Recall that $K_{-2}^{\text{MW}}(F)$ is generated by the $\langle a \rangle \eta^2$ with $a \in F^*$.

The Hopf link

Recall that we fixed coordinates x, y, z, t for \mathbb{A}_F^4 and u, v for \mathbb{A}_F^2 .

- The image of the Hopf link:

$$\{x = 0, y = 0\} \sqcup \{z = 0, t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

- The parametrization of the Hopf link:

$$\varphi_1 : (x, y, z, t) \leftrightarrow (0, 0, u, v), \varphi_2 : (x, y, z, t) \leftrightarrow (u, v, 0, 0)$$

- The orientation of the Hopf link:

$$\sigma_1 : \bar{x}^* \wedge \bar{y}^* \mapsto 1, \sigma_2 : \bar{z}^* \wedge \bar{t}^* \mapsto 1$$

The quadratic linking degree of the Hopf link

Or. fund. classes	$\eta \otimes (\bar{x}^* \wedge \bar{y}^*)$		$\eta \otimes (\bar{z}^* \wedge \bar{t}^*)$
Seifert classes	$\langle x \rangle \otimes \bar{y}^*$		$\langle z \rangle \otimes \bar{t}^*$
Apply int. prod.	$\langle xz \rangle \otimes (\bar{t}^* \wedge \bar{y}^*)$		
Quad. link. class	$-\langle z \rangle \eta \otimes (\bar{t}^* \wedge \bar{x}^* \wedge \bar{y}^*)$	\oplus	$\langle x \rangle \eta \otimes (\bar{y}^* \wedge \bar{z}^* \wedge \bar{t}^*)$
Apply $\tilde{o}_1 \oplus \tilde{o}_2$	$-\langle z \rangle \eta \otimes \bar{t}^*$	\oplus	$\langle x \rangle \eta \otimes \bar{y}^*$
Apply $\varphi_1^* \oplus \varphi_2^*$	$-\langle u \rangle \eta \otimes \bar{v}^*$	\oplus	$\langle u \rangle \eta \otimes \bar{v}^*$
Apply $\partial \oplus \partial$	$-\eta^2 \otimes (\bar{u}^* \wedge \bar{v}^*)$	\oplus	$\eta^2 \otimes (\bar{u}^* \wedge \bar{v}^*)$
Quad. link. degree	$-\eta^2$	\oplus	η^2

A variant of the Hopf link

- The image is the same as the Hopf link's image:

$$\{x = y, y = 0\} \sqcup \{z = 0, a \times t = 0\} \subset \mathbb{A}_F^4 \setminus \{0\} \text{ with } a \in F^*$$

- The parametrization is the same:

$$\varphi_1 : (x, y, z, t) \leftrightarrow (0, 0, u, v), \varphi_2 : (x, y, z, t) \leftrightarrow (u, v, 0, 0)$$

- The orientation is different:

$$\alpha_1 : \overline{x - y}^* \wedge \overline{y}^* \mapsto 1, \alpha_2 : \overline{z}^* \wedge \overline{at}^* \mapsto 1$$

The quadratic linking degree of a variant of the Hopf link

$$[o_1^{var}] = \eta \otimes \overline{x-y}^* \wedge \overline{y}^* = [o_1^{Hopf}] \quad [o_2^{var}] = \eta \otimes \overline{z}^* \wedge \overline{at}^* = \langle a \rangle [o_2^{Hopf}]$$

$$\text{since } \begin{pmatrix} x-y \\ y \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{since } \begin{pmatrix} z \\ at \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix} \begin{pmatrix} z \\ t \end{pmatrix}$$

$$\mathcal{S}_1^{var} = \mathcal{S}_1^{Hopf}$$

$$\mathcal{S}_2^{var} = \langle a \rangle \mathcal{S}_2^{Hopf}$$

$$\mathcal{S}_1^{var} \cdot \mathcal{S}_2^{var} = \langle a \rangle \mathcal{S}_1^{Hopf} \cdot \mathcal{S}_2^{Hopf}$$

$$\partial(\mathcal{S}_1^{var} \cdot \mathcal{S}_2^{var}) = \langle a \rangle \partial(\mathcal{S}_1^{Hopf} \cdot \mathcal{S}_2^{Hopf})$$

The quadratic linking degree is $(-\langle a \rangle \eta^2, \eta^2)$.

Fact

Let \mathcal{L} be an oriented link with two components of quadratic linking degree $(d_1, d_2) \in K_{-2}^{\text{MW}}(F) \oplus K_{-2}^{\text{MW}}(F)$. Let $a = (a_1, a_2)$ be a couple of elements of F^* and \mathcal{L}_a be the link obtained from \mathcal{L} by changing the orientation o_1 into $o_1 \circ (\times a_1)$ and the orientation o_2 into $o_2 \circ (\times a_2)$. Then $\text{Qlc}_{\mathcal{L}_a} = \langle a_1 a_2 \rangle \text{Qlc}_{\mathcal{L}}$ and $\text{Qld}_{\mathcal{L}_a} = (\langle a_2 \rangle d_1, \langle a_1 \rangle d_2)$.

Fact

Let \mathcal{L} be an oriented link with two components of quadratic linking degree $(d_1, d_2) \in K_{-2}^{\text{MW}}(F) \oplus K_{-2}^{\text{MW}}(F)$. Let $a = (a_1, a_2)$ be a couple of elements of F^* and \mathcal{L}_a be the link obtained from \mathcal{L} by changing the orientation o_1 into $o_1 \circ (\times a_1)$ and the orientation o_2 into $o_2 \circ (\times a_2)$. Then $\text{Qlc}_{\mathcal{L}_a} = \langle a_1 a_2 \rangle \text{Qlc}_{\mathcal{L}}$ and $\text{Qld}_{\mathcal{L}_a} = (\langle a_2 \rangle d_1, \langle a_1 \rangle d_2)$.

Similarly, changes of parametrizations of the link can only multiply each component of the quadratic linking degree by elements of the form $\langle a \rangle$ with $a \in F^*$ (and do not change the quadratic linking class).

Fact

Let \mathcal{L} be an oriented link with two components of quadratic linking degree $(d_1, d_2) \in K_{-2}^{\text{MW}}(F) \oplus K_{-2}^{\text{MW}}(F)$. Let $a = (a_1, a_2)$ be a couple of elements of F^* and \mathcal{L}_a be the link obtained from \mathcal{L} by changing the orientation o_1 into $o_1 \circ (\times a_1)$ and the orientation o_2 into $o_2 \circ (\times a_2)$. Then $\text{Qlc}_{\mathcal{L}_a} = \langle a_1 a_2 \rangle \text{Qlc}_{\mathcal{L}}$ and $\text{Qld}_{\mathcal{L}_a} = (\langle a_2 \rangle d_1, \langle a_1 \rangle d_2)$.

Similarly, changes of parametrizations of the link can only multiply each component of the quadratic linking degree by elements of the form $\langle a \rangle$ with $a \in F^*$ (and do not change the quadratic linking class).

We want invariants of the quadratic linking degree. (Similarly to the absolute value of the linking number in knot theory)

Why a “quadratic” linking degree?

- The (commutative) ring with unit $K_0^{\text{MW}}(F)$ is isomorphic to the Grothendieck-Witt ring $\text{GW}(F)$ of F via $\langle a \rangle \in K_0^{\text{MW}}(F) \leftrightarrow \langle a \rangle \in \text{GW}(F)$.
- For all $n < 0$, the abelian group $K_n^{\text{MW}}(F)$ is isomorphic to the Witt group $W(F)$ of F via $\langle a \rangle \eta^{-n} \in K_n^{\text{MW}}(F) \leftrightarrow \langle a \rangle \in W(F)$.

The real definition of the quadratic linking degree

We define the quadratic linking degree of \mathcal{L} as the image of the quadratic linking class of \mathcal{L} by the isomorphism

$$H^1(Z, \underline{K}_0^{\text{MW}}\{\nu_Z\}) \rightarrow K_{-2}^{\text{MW}}(F) \oplus K_{-2}^{\text{MW}}(F) \rightarrow W(F) \oplus W(F).$$

Why a “quadratic” linking degree?

- The (commutative) ring with unit $K_0^{\text{MW}}(F)$ is isomorphic to the Grothendieck-Witt ring $\text{GW}(F)$ of F via $\langle a \rangle \in K_0^{\text{MW}}(F) \leftrightarrow \langle a \rangle \in \text{GW}(F)$.
- For all $n < 0$, the abelian group $K_n^{\text{MW}}(F)$ is isomorphic to the Witt group $W(F)$ of F via $\langle a \rangle \eta^{-n} \in K_n^{\text{MW}}(F) \leftrightarrow \langle a \rangle \in W(F)$.

The real definition of the quadratic linking degree

We define the quadratic linking degree of \mathcal{L} as the image of the quadratic linking class of \mathcal{L} by the isomorphism

$$H^1(Z, \underline{K}_0^{\text{MW}}\{\nu_Z\}) \rightarrow K_{-2}^{\text{MW}}(F) \oplus K_{-2}^{\text{MW}}(F) \rightarrow W(F) \oplus W(F).$$

The Grothendieck-Witt ring of F and the Witt ring of F (and underlying Witt group of F) are constructed from symmetric bilinear forms on F . If F is of characteristic different from 2 (i.e. $2 \neq 0$ in F) then they are also constructed from quadratic forms.

Interlude: symmetric bilinear forms and quadratic forms

Definition

- A bilinear form on an F -vector space V of finite dimension is a bilinear map $b : V \times V \rightarrow F$. It is symmetric if for all $v, w \in V$, $b(v, w) = b(w, v)$.
- If F is of characteristic different from 2, a quadratic form on V is a map $q : V \rightarrow F$ such that the map $b : \begin{cases} V \times V & \rightarrow & F \\ (x, y) & \mapsto & \frac{1}{2}(q(x+y) - q(x) - q(y)) \end{cases}$ is a symmetric bilinear form such that for all $x \in V$, $b(x, x) = q(x)$. We call b the polar form of q .

Note that if $b : V \times V \rightarrow F$ is a symmetric bilinear form and F is of characteristic different from 2 then $q : \begin{cases} V & \rightarrow & F \\ x & \mapsto & b(x, x) \end{cases}$ is a quadratic form (of polar form b).

Definition

Let $b : V \times V \rightarrow F$ and $b' : V' \times V' \rightarrow F$ be symmetric bilinear forms.

- The (orthogonal) sum of b and b' is the symmetric bilinear form $b \perp b' : (V \oplus V') \times (V \oplus V') \rightarrow F$ which sends $((x, x'), (y, y'))$ to $b(x, y) + b'(x', y')$.
- The (tensor) product of b and b' is the symmetric bilinear form $b \otimes b' : (V \otimes V') \times (V \otimes V') \rightarrow F$ which sends $(\sum_{i \in I} x_i \otimes x'_i, \sum_{j \in J} y_j \otimes y'_j)$ to $\sum_{(i,j) \in I \times J} b(x_i, y_j) \times b'(x'_i, y'_j)$.

Definition

- The symmetric bilinear form $b : V \times V \rightarrow F$ is non-degenerate if 0 is the only element x of V which verifies that for all $y \in V$, $b(x, y) = 0$.
- Two non-degenerate symmetric bilinear forms $b : V \times V \rightarrow F$ and $b' : V' \times V' \rightarrow F$ are isometric if there exists a linear isomorphism $u : V \rightarrow V'$ such that for all $x, y \in V$, $b(x, y) = b'(u(x), u(y))$.

This gives a structure of commutative semiring (commutative monoid + commutative product) on the isometry classes. Grothendieck's construction gives a commutative ring: the Grothendieck-Witt ring of F . Its elements are \mathbb{Z} -linear combinations of the classes

$$\langle a \rangle : \begin{cases} F \times F & \rightarrow & F \\ (x, y) & \mapsto & axy \end{cases} \text{ of symmetric bilinear forms (with } a \in F^* \text{)}.$$

If F is of characteristic $\neq 2$, as a quadratic form $\langle a \rangle : \begin{cases} F & \rightarrow & F \\ x & \mapsto & ax^2 \end{cases}$.

Definition

- The hyperbolic plane $b_h : F^2 \times F^2 \rightarrow F$ is the symmetric bilinear form which sends $((x_1, y_1), (x_2, y_2))$ to $x_1y_2 + x_2y_1$.
- Two non-degenerate symmetric bilinear forms $b : V \times V \rightarrow F$ and $b' : V' \times V' \rightarrow F$ are Witt-equivalent if there exist $m, n \geq 0$ integers such that $b \perp mb_h$ is isometric to $b' \perp nb_h$.

This gives a structure of commutative ring on the Witt-equivalence classes: the Witt ring of F . Its elements are sums of classes

$$\langle a \rangle : \begin{cases} F \times F & \rightarrow & F \\ (x, y) & \mapsto & axy \end{cases} \text{ of symmetric bilinear forms (with } a \in F^* \text{)}.$$

If F is of characteristic $\neq 2$, as a quadratic form $\langle a \rangle : \begin{cases} F & \rightarrow & F \\ x & \mapsto & ax^2. \end{cases}$

Presentations of $\mathrm{GW}(F)$ and $W(F)$

- As a commutative ring (resp. abelian group), the Grothendieck-Witt ring (resp. group) $\mathrm{GW}(F)$ is generated by the $\langle a \rangle$ for $a \in F^*$ subject to the relations :
 - $\langle ab^2 \rangle = \langle a \rangle$ for all $a, b \in F^*$;
 - $\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \langle (a + b)ab \rangle$ for all $a, b \in F^*$ such that $a + b \in F^*$.

Presentations of $GW(F)$ and $W(F)$

- As a commutative ring (resp. abelian group), the Grothendieck-Witt ring (resp. group) $GW(F)$ is generated by the $\langle a \rangle$ for $a \in F^*$ subject to the relations :
 - $\langle ab^2 \rangle = \langle a \rangle$ for all $a, b \in F^*$;
 - $\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \langle (a + b)ab \rangle$ for all $a, b \in F^*$ such that $a + b \in F^*$.
- As a commutative ring (resp. abelian group), the Witt ring (resp. group) $W(F)$ is generated by the $\langle a \rangle$ for $a \in F^*$ subject to the relations :
 - $\langle ab^2 \rangle = \langle a \rangle$ for all $a, b \in F^*$;
 - $\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \langle (a + b)ab \rangle$ for all $a, b \in F^*$ such that $a + b \in F^*$;
 - $\langle 1 \rangle + \langle -1 \rangle = 0$.

Presentations of $GW(F)$ and $W(F)$

- As a commutative ring (resp. abelian group), the Grothendieck-Witt ring (resp. group) $GW(F)$ is generated by the $\langle a \rangle$ for $a \in F^*$ subject to the relations :
 - $\langle ab^2 \rangle = \langle a \rangle$ for all $a, b \in F^*$;
 - $\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \langle (a + b)ab \rangle$ for all $a, b \in F^*$ such that $a + b \in F^*$.
- As a commutative ring (resp. abelian group), the Witt ring (resp. group) $W(F)$ is generated by the $\langle a \rangle$ for $a \in F^*$ subject to the relations :
 - $\langle ab^2 \rangle = \langle a \rangle$ for all $a, b \in F^*$;
 - $\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \langle (a + b)ab \rangle$ for all $a, b \in F^*$ such that $a + b \in F^*$;
 - $\langle 1 \rangle + \langle -1 \rangle = 0$.
- This last relation corresponds to the vanishing of the hyperbolic plane.

Examples of Witt rings

- $W(\mathbb{C}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2 (where the rank of $\sum_{i=1}^n \langle a_i \rangle$ is n)

Examples of Witt rings

- $W(\mathbb{C}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2 (where the rank of $\sum_{i=1}^n \langle a_i \rangle$ is n)
- $W(\mathbb{R}) \simeq \mathbb{Z}$ via the signature (where the signature of $\sum_{i=1}^p \langle 1 \rangle + \sum_{j=1}^q \langle -1 \rangle$ is $p - q$)

Examples of Witt rings

- $W(\mathbb{C}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2 (where the rank of $\sum_{i=1}^n \langle a_i \rangle$ is n)
- $W(\mathbb{R}) \simeq \mathbb{Z}$ via the signature (where the signature of $\sum_{i=1}^p \langle 1 \rangle + \sum_{j=1}^q \langle -1 \rangle$ is $p - q$)
- $W(\mathbb{Q}) \simeq W(\mathbb{R}) \oplus \bigoplus_{p \text{ prime}} W(\mathbb{Z}/p\mathbb{Z})$ via $\langle a \rangle \mapsto \langle a \rangle + \sum_{p \text{ prime divisor of } a} \langle \frac{a}{p} \rangle$

Examples of Witt rings

- $W(\mathbb{C}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2 (where the rank of $\sum_{i=1}^n \langle a_i \rangle$ is n)
- $W(\mathbb{R}) \simeq \mathbb{Z}$ via the signature (where the signature of $\sum_{i=1}^p \langle 1 \rangle + \sum_{j=1}^q \langle -1 \rangle$ is $p - q$)
- $W(\mathbb{Q}) \simeq W(\mathbb{R}) \oplus \bigoplus_{p \text{ prime}} W(\mathbb{Z}/p\mathbb{Z})$ via $\langle a \rangle \mapsto \langle a \rangle + \sum_{p \text{ prime divisor of } a} \langle \frac{a}{p} \rangle$
- $W(\mathbb{Z}/2\mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2

Examples of Witt rings

- $W(\mathbb{C}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2 (where the rank of $\sum_{i=1}^n \langle a_i \rangle$ is n)
- $W(\mathbb{R}) \simeq \mathbb{Z}$ via the signature (where the signature of $\sum_{i=1}^p \langle 1 \rangle + \sum_{j=1}^q \langle -1 \rangle$ is $p - q$)
- $W(\mathbb{Q}) \simeq W(\mathbb{R}) \oplus \bigoplus_{p \text{ prime}} W(\mathbb{Z}/p\mathbb{Z})$ via $\langle a \rangle \mapsto \langle a \rangle + \sum_{p \text{ prime divisor of } a} \langle \frac{a}{p} \rangle$
- $W(\mathbb{Z}/2\mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2
- For all $p \equiv 3 \pmod{4}$, $W(\mathbb{Z}/p\mathbb{Z}) \simeq \mathbb{Z}/4\mathbb{Z}$ via the signature modulo 4

Examples of Witt rings

- $W(\mathbb{C}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2 (where the rank of $\sum_{i=1}^n \langle a_i \rangle$ is n)
- $W(\mathbb{R}) \simeq \mathbb{Z}$ via the signature (where the signature of $\sum_{i=1}^p \langle 1 \rangle + \sum_{j=1}^q \langle -1 \rangle$ is $p - q$)
- $W(\mathbb{Q}) \simeq W(\mathbb{R}) \oplus \bigoplus_{p \text{ prime}} W(\mathbb{Z}/p\mathbb{Z})$ via $\langle a \rangle \mapsto \langle a \rangle + \sum_{p \text{ prime divisor of } a} \langle \frac{a}{p} \rangle$
- $W(\mathbb{Z}/2\mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$ via the rank modulo 2
- For all $p \equiv 3 \pmod{4}$, $W(\mathbb{Z}/p\mathbb{Z}) \simeq \mathbb{Z}/4\mathbb{Z}$ via the signature modulo 4
- For all $p \equiv 1 \pmod{4}$, $W(\mathbb{Z}/p\mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ via the “signature couple” modulo 2 (if $a \in \mathbb{Z}/p\mathbb{Z}$ is not a square, $\sum_{i=1}^p \langle 1 \rangle + \sum_{j=1}^q \langle a \rangle$ is sent to $(p \pmod{2}, q \pmod{2})$)

It is difficult in general to know if two elements of the Witt group $W(F)$ are equal. For instance, let $a, b, c, d \in F^*$ such that d is not a square in F^* and such that (1) and (2) below are well-defined. Can you tell which of the two following elements of $W(F)$ is equal to $\langle a \rangle + \langle b \rangle$? (There is exactly one which is equal to $\langle a \rangle + \langle b \rangle$)

① $\langle (a + b)c^2 + (a + b)abd \rangle + \langle (a + b)(c^2 + abd)abd \rangle$

② $\langle (a + b)c^2 + (a + b)abd^2 \rangle + \langle (a + b)(c^2 + abd^2)ab \rangle$

Recall that the relations in $W(F)$ are:

- $\langle ab^2 \rangle = \langle a \rangle$ for all $a, b \in F^*$;
- $\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \langle (a + b)ab \rangle$ for all $a, b \in F^*$ such that $a + b \in F^*$;
- $\langle 1 \rangle + \langle -1 \rangle = 0$.

Solution

The second one is equal to $\langle a \rangle + \langle b \rangle$. Indeed:

$$\begin{aligned}\langle a \rangle + \langle b \rangle &= \langle (a+b)c^2 \rangle + \langle (a+b)abd^2 \rangle \\ &= \langle (a+b)c^2 + (a+b)abd^2 \rangle + \langle (a+b)(c^2 + abd^2)ab(a+b)^2c^2d^2 \rangle \\ &= \langle (a+b)c^2 + (a+b)abd^2 \rangle + \langle (a+b)(c^2 + abd^2)ab \rangle\end{aligned}$$

To see that the first one is different from $\langle a \rangle + \langle b \rangle$, we will use one of the invariants presented later in this talk.

Invariants by multiplication by $\langle a \rangle$ for all $a \in F^*$

Case $F = \mathbb{R}$

If $F = \mathbb{R}$, the absolute value of an element of $W(\mathbb{R}) \simeq \mathbb{Z}$ is invariant by multiplication by $\langle a \rangle$ for all $a \in F^*$.

Invariants by multiplication by $\langle a \rangle$ for all $a \in F^*$

Case $F = \mathbb{R}$

If $F = \mathbb{R}$, the absolute value of an element of $W(\mathbb{R}) \simeq \mathbb{Z}$ is invariant by multiplication by $\langle a \rangle$ for all $a \in F^*$.

General case

The rank modulo 2 is invariant by multiplication by $\langle a \rangle$ for all $a \in F^*$.

Definition

Let $d = \sum_{i=1}^n \langle a_i \rangle \in W(F)$. There exists a unique sequence of abelian groups $Q_{d,k}$ and of elements $\Sigma_k(d) \in Q_{d,k}$, where k ranges over the nonnegative even integers, such that:

- $Q_{d,0} = W(F)$ and $\Sigma_0(d) = 1 \in Q_{d,0}$;
- for each positive even integer k , $Q_{d,k}$ is the quotient group $Q_{d,k-2}/(\Sigma_{k-2}(d))$;

- for each positive even integer k ,
$$\Sigma_k(d) = \sum_{1 \leq i_1 < \dots < i_k \leq n} \langle \prod_{1 \leq j \leq k} a_{i_j} \rangle \in Q_{d,k}.$$

General case

The Σ_k are invariant by multiplication by $\langle a \rangle$ for all $a \in F^*$.

- $\Sigma_2 : \begin{cases} W(F) & \rightarrow W(F)/(1) \\ \sum_{i=1}^n \langle a_i \rangle & \mapsto \sum_{1 \leq i < j \leq n} \langle a_i a_j \rangle \end{cases}$ (if $n < 2$, it sends $\sum_{i=1}^n \langle a_i \rangle$ to 0)

- $\Sigma_2 : \begin{cases} W(F) & \rightarrow W(F)/(1) \\ \sum_{i=1}^n \langle a_i \rangle & \mapsto \sum_{1 \leq i < j \leq n} \langle a_i a_j \rangle \end{cases}$ (if $n < 2$, it sends $\sum_{i=1}^n \langle a_i \rangle$ to 0)
- This is not interesting if $W(F)/(1) = 0$ (for instance if $F = \mathbb{R}$).

- $\Sigma_2 : \begin{cases} W(F) & \rightarrow W(F)/(1) \\ \sum_{i=1}^n \langle a_i \rangle & \mapsto \sum_{1 \leq i < j \leq n} \langle a_i a_j \rangle \end{cases}$ (if $n < 2$, it sends $\sum_{i=1}^n \langle a_i \rangle$ to 0)
- This is not interesting if $W(F)/(1) = 0$ (for instance if $F = \mathbb{R}$).
- It is interesting for $F = \mathbb{Q}$ for instance: $W(\mathbb{Q})/(1) \simeq \bigoplus_{p \text{ prime}} W(\mathbb{Z}/p\mathbb{Z})$.

- $\Sigma_2 : \begin{cases} W(F) & \rightarrow W(F)/(1) \\ \sum_{i=1}^n \langle a_i \rangle & \mapsto \sum_{1 \leq i < j \leq n} \langle a_i a_j \rangle \end{cases}$ (if $n < 2$, it sends $\sum_{i=1}^n \langle a_i \rangle$ to 0)
- This is not interesting if $W(F)/(1) = 0$ (for instance if $F = \mathbb{R}$).
- It is interesting for $F = \mathbb{Q}$ for instance: $W(\mathbb{Q})/(1) \simeq \bigoplus_{p \text{ prime}} W(\mathbb{Z}/p\mathbb{Z})$.

- $\Sigma_4 : \begin{cases} W(F) & \rightarrow \bigcup_{d \in W(F)} (W(F)/(1))/(\Sigma_2(d)) \\ \sum_{i=1}^n \langle a_i \rangle & \mapsto \sum_{1 \leq i < j < k < l \leq n} \langle a_i a_j a_k a_l \rangle \end{cases}$

- $\Sigma_2 : \begin{cases} W(F) & \rightarrow W(F)/(1) \\ \sum_{i=1}^n \langle a_i \rangle & \mapsto \sum_{1 \leq i < j \leq n} \langle a_i a_j \rangle \end{cases}$ (if $n < 2$, it sends $\sum_{i=1}^n \langle a_i \rangle$ to 0)
- This is not interesting if $W(F)/(1) = 0$ (for instance if $F = \mathbb{R}$).
- It is interesting for $F = \mathbb{Q}$ for instance: $W(\mathbb{Q})/(1) \simeq \bigoplus_{p \text{ prime}} W(\mathbb{Z}/p\mathbb{Z})$.

- $\Sigma_4 : \begin{cases} W(F) & \rightarrow \bigcup_{d \in W(F)} (W(F)/(1))/(\Sigma_2(d)) \\ \sum_{i=1}^n \langle a_i \rangle & \mapsto \sum_{1 \leq i < j < k < l \leq n} \langle a_i a_j a_k a_l \rangle \end{cases}$

- We only want to compare $\Sigma_4(d)$ and $\Sigma_4(d')$ if $\Sigma_2(d) = \Sigma_2(d')$.

$\Sigma_2(\langle (a+b)c^2 + (a+b)abd \rangle + \langle (a+b)(c^2 + abd)abd \rangle) =$
 $\langle ((a+b)c^2 + (a+b)abd)(a+b)(c^2 + abd)abd \rangle = \langle abd \rangle \neq \langle ab \rangle \in W(F)/(1)$
 since d is not a square in F^* . Since $\Sigma_2(\langle a \rangle + \langle b \rangle) = \langle ab \rangle$,

$$\langle a \rangle + \langle b \rangle \neq \langle (a+b)c^2 + (a+b)abd \rangle + \langle (a+b)(c^2 + abd)abd \rangle \in W(F)$$

Application: invariants of the quadratic linking degree

Let \mathcal{L} be an oriented link with two components (in motivic knot theory). We denote by $(d_1, d_2) \in W(F) \oplus W(F)$ its quadratic linking degree.

- If $F = \mathbb{R}$ then the absolute value of d_1 and the absolute value of d_2 are invariant under changes of orientations σ_1, σ_2 and of parametrizations of $\varphi_1, \varphi_2 : \mathbb{A}_{\mathbb{R}}^2 \setminus \{0\} \rightarrow \mathbb{A}_{\mathbb{R}}^4 \setminus \{0\}$.

Application: invariants of the quadratic linking degree

Let \mathcal{L} be an oriented link with two components (in motivic knot theory). We denote by $(d_1, d_2) \in W(F) \oplus W(F)$ its quadratic linking degree.

- If $F = \mathbb{R}$ then the absolute value of d_1 and the absolute value of d_2 are invariant under changes of orientations σ_1, σ_2 and of parametrizations of $\varphi_1, \varphi_2 : \mathbb{A}_{\mathbb{R}}^2 \setminus \{0\} \rightarrow \mathbb{A}_{\mathbb{R}}^4 \setminus \{0\}$.
- The rank modulo 2 of d_1 and the rank modulo 2 of d_2 are invariant under changes of orientations σ_1, σ_2 and of parametrizations of $\varphi_1, \varphi_2 : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$.

Application: invariants of the quadratic linking degree

Let \mathcal{L} be an oriented link with two components (in motivic knot theory). We denote by $(d_1, d_2) \in W(F) \oplus W(F)$ its quadratic linking degree.

- If $F = \mathbb{R}$ then the absolute value of d_1 and the absolute value of d_2 are invariant under changes of orientations σ_1, σ_2 and of parametrizations of $\varphi_1, \varphi_2 : \mathbb{A}_{\mathbb{R}}^2 \setminus \{0\} \rightarrow \mathbb{A}_{\mathbb{R}}^4 \setminus \{0\}$.
- The rank modulo 2 of d_1 and the rank modulo 2 of d_2 are invariant under changes of orientations σ_1, σ_2 and of parametrizations of $\varphi_1, \varphi_2 : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$.
- For every positive even integer k , $\Sigma_k(d_1)$ and $\Sigma_k(d_2)$ are invariant under changes of orientations σ_1, σ_2 and of parametrizations of $\varphi_1, \varphi_2 : \mathbb{A}_F^2 \setminus \{0\} \rightarrow \mathbb{A}_F^4 \setminus \{0\}$.

Another Hopf link

From now on, F is a perfect field of characteristic different from 2. Recall that we fixed coordinates x, y, z, t for \mathbb{A}_F^4 and u, v for \mathbb{A}_F^2 .

- The image is different from the Hopf link we saw before:

$$\{z = x, t = y\} \sqcup \{z = -x, t = -y\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

But the change of coordinates $x' = z - x$, $y' = t - y$, $z' = z + x$, $t' = t + y$ would give $\{x' = 0, y' = 0\} \sqcup \{z' = 0, t' = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$.

Another Hopf link

From now on, F is a perfect field of characteristic different from 2. Recall that we fixed coordinates x, y, z, t for \mathbb{A}_F^4 and u, v for \mathbb{A}_F^2 .

- The image is different from the Hopf link we saw before:

$$\{z = x, t = y\} \sqcup \{z = -x, t = -y\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

But the change of coordinates $x' = z - x$, $y' = t - y$, $z' = z + x$, $t' = t + y$ would give $\{x' = 0, y' = 0\} \sqcup \{z' = 0, t' = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$.

- The parametrization is $\varphi_1 : (x, y, z, t) \leftrightarrow (u, v, u, v)$ and $\varphi_2 : (x, y, z, t) \leftrightarrow (u, v, -u, -v)$.

Another Hopf link

From now on, F is a perfect field of characteristic different from 2. Recall that we fixed coordinates x, y, z, t for \mathbb{A}_F^4 and u, v for \mathbb{A}_F^2 .

- The image is different from the Hopf link we saw before:

$$\{z = x, t = y\} \sqcup \{z = -x, t = -y\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

But the change of coordinates $x' = z - x$, $y' = t - y$, $z' = z + x$, $t' = t + y$ would give $\{x' = 0, y' = 0\} \sqcup \{z' = 0, t' = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$.

- The parametrization is $\varphi_1 : (x, y, z, t) \leftrightarrow (u, v, u, v)$ and $\varphi_2 : (x, y, z, t) \leftrightarrow (u, v, -u, -v)$.
- The orientation is the following:

$$\omega_1 : \overline{z - x}^* \wedge \overline{t - y}^* \mapsto 1, \omega_2 : \overline{z + x}^* \wedge \overline{t + y}^* \mapsto 1$$

- This Hopf link is an analogue of the Hopf link in knot theory! In knot theory, the Hopf link is given by $\{z = x, t = y\} \sqcup \{z = -x, t = -y\}$ in $\mathbb{S}_\varepsilon^3 = \{(x, y, z, t) \in \mathbb{R}^4, x^2 + y^2 + z^2 + t^2 = \varepsilon^2\}$ for ε small enough and has linking number 1.

- This Hopf link is an analogue of the Hopf link in knot theory! In knot theory, the Hopf link is given by $\{z = x, t = y\} \sqcup \{z = -x, t = -y\}$ in $\mathbb{S}_\varepsilon^3 = \{(x, y, z, t) \in \mathbb{R}^4, x^2 + y^2 + z^2 + t^2 = \varepsilon^2\}$ for ε small enough and has linking number 1.
- Its quadratic linking degree is $(\langle 1 \rangle, \langle -1 \rangle) = (1, -1) \in W(F) \oplus W(F)$.

- This Hopf link is an analogue of the Hopf link in knot theory! In knot theory, the Hopf link is given by $\{z = x, t = y\} \sqcup \{z = -x, t = -y\}$ in $\mathbb{S}_\varepsilon^3 = \{(x, y, z, t) \in \mathbb{R}^4, x^2 + y^2 + z^2 + t^2 = \varepsilon^2\}$ for ε small enough and has linking number 1.
- Its quadratic linking degree is $(\langle 1 \rangle, \langle -1 \rangle) = (1, -1) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle, \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.

- This Hopf link is an analogue of the Hopf link in knot theory! In knot theory, the Hopf link is given by $\{z = x, t = y\} \sqcup \{z = -x, t = -y\}$ in $\mathbb{S}_\varepsilon^3 = \{(x, y, z, t) \in \mathbb{R}^4, x^2 + y^2 + z^2 + t^2 = \varepsilon^2\}$ for ε small enough and has linking number 1.
- Its quadratic linking degree is $(\langle 1 \rangle, \langle -1 \rangle) = (1, -1) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle, \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.
- If $F = \mathbb{R}$, the absolute value of each component is 1.

- This Hopf link is an analogue of the Hopf link in knot theory! In knot theory, the Hopf link is given by $\{z = x, t = y\} \sqcup \{z = -x, t = -y\}$ in $\mathbb{S}_\varepsilon^3 = \{(x, y, z, t) \in \mathbb{R}^4, x^2 + y^2 + z^2 + t^2 = \varepsilon^2\}$ for ε small enough and has linking number 1.
- Its quadratic linking degree is $(\langle 1 \rangle, \langle -1 \rangle) = (1, -1) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle, \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.
- If $F = \mathbb{R}$, the absolute value of each component is 1.
- The rank modulo 2 of each component is 1.

- This Hopf link is an analogue of the Hopf link in knot theory! In knot theory, the Hopf link is given by $\{z = x, t = y\} \sqcup \{z = -x, t = -y\}$ in $\mathbb{S}_\varepsilon^3 = \{(x, y, z, t) \in \mathbb{R}^4, x^2 + y^2 + z^2 + t^2 = \varepsilon^2\}$ for ε small enough and has linking number 1.
- Its quadratic linking degree is $(\langle 1 \rangle, \langle -1 \rangle) = (1, -1) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle, \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.
- If $F = \mathbb{R}$, the absolute value of each component is 1.
- The rank modulo 2 of each component is 1.
- For every positive even integer k , the image by Σ_k of each component is 0.

The Solomon link

- In knot theory, the Solomon link is given by $\{z = x^2 - y^2, t = 2xy\} \sqcup \{z = -x^2 + y^2, t = -2xy\}$ in \mathbb{S}_ε^3 for ε small enough and has linking number 2.
- In motivic knot theory, the image of the Solomon link is:

$$\{z = x^2 - y^2, t = 2xy\} \sqcup \{z = -x^2 + y^2, t = -2xy\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

- The parametrization is $\varphi_1 : (x, y, z, t) \leftrightarrow (u, v, u^2 - v^2, 2uv)$ and $\varphi_2 : (x, y, z, t) \leftrightarrow (u, v, -u^2 + v^2, -2uv)$.
- The orientation is the following:

$$o_1 : \overline{z - x^2 + y^2}^* \wedge \overline{t - 2xy}^* \mapsto 1, o_2 : \overline{z + x^2 - y^2}^* \wedge \overline{t + 2xy}^* \mapsto 1$$

- Its quadratic linking degree is $(\langle 1 \rangle + \langle 1 \rangle, \langle -1 \rangle + \langle -1 \rangle) = (2, -2) \in W(F) \oplus W(F)$.

- Its quadratic linking degree is $(\langle 1 \rangle + \langle 1 \rangle, \langle -1 \rangle + \langle -1 \rangle) = (2, -2) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle a \rangle, \langle b \rangle + \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.

- Its quadratic linking degree is $(\langle 1 \rangle + \langle 1 \rangle, \langle -1 \rangle + \langle -1 \rangle) = (2, -2) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle a \rangle, \langle b \rangle + \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.
- If $F = \mathbb{R}$, the absolute value of each component is 2.

- Its quadratic linking degree is $(\langle 1 \rangle + \langle 1 \rangle, \langle -1 \rangle + \langle -1 \rangle) = (2, -2) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle a \rangle, \langle b \rangle + \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.
- If $F = \mathbb{R}$, the absolute value of each component is 2.
- The rank modulo 2 of each component is 0.

- Its quadratic linking degree is $(\langle 1 \rangle + \langle 1 \rangle, \langle -1 \rangle + \langle -1 \rangle) = (2, -2) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle a \rangle, \langle b \rangle + \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.
- If $F = \mathbb{R}$, the absolute value of each component is 2.
- The rank modulo 2 of each component is 0.
- For every positive even integer k , the image by Σ_k of each component is 0.

- Its quadratic linking degree is $(\langle 1 \rangle + \langle 1 \rangle, \langle -1 \rangle + \langle -1 \rangle) = (2, -2) \in W(F) \oplus W(F)$.
- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle a \rangle, \langle b \rangle + \langle b \rangle) \in W(F) \oplus W(F)$ with $a, b \in F^*$.
- If $F = \mathbb{R}$, the absolute value of each component is 2.
- The rank modulo 2 of each component is 0.
- For every positive even integer k , the image by Σ_k of each component is 0.
- More generally, we have analogues of the torus links $T(2, 2n)$ (of linking number n); the quadratic linking degree of $T(2, 2n)$ is $(n, -n) \in W(F) \oplus W(F)$, which gives n as absolute value if $F = \mathbb{R}$, n modulo 2 as rank modulo 2, and 0 for the Σ_k .

Binary links

- The image of the binary link B_a with $a \in F^* \setminus \{-1\}$:

$$\{f_1 = 0, g_1 = 0\} \sqcup \{f_2 = 0, g_2 = 0\} \subset \mathbb{A}_F^4 \setminus \{0\}$$

with $f_1 = t - ((1 + a)x - y)y$, $g_1 = z - x(x - y)$,
 $f_2 = t + ((1 + a)x - y)y$, $g_2 = z + x(x - y)$.

- The parametrization of the binary link B_a :

$$\varphi_1 : (x, y, z, t) \leftrightarrow (u, v, ((1 + a)u - v)v, u(u - v))$$

$$\varphi_2 : (x, y, z, t) \leftrightarrow (u, v, -((1 + a)u - v)v, -u(u - v))$$

- The orientation of the binary link B_a :

$$\alpha_1 : \overline{f_1}^* \wedge \overline{g_1}^* \mapsto 1, \alpha_2 : \overline{f_2}^* \wedge \overline{g_2}^* \mapsto 1$$

Or. fund. cyc.	$\eta \otimes (\overline{f_1}^* \wedge \overline{g_1}^*)$		$\eta \otimes (\overline{f_2}^* \wedge \overline{g_2}^*)$
Seifert divisors	$\langle f_1 \rangle \otimes \overline{g_1}^*$		$\langle f_2 \rangle \otimes \overline{g_2}^*$
Apply inter. prod.	$\langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x - y)$ $+ \langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x)$		
...	...		
Apply $\partial \oplus \partial$	$(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$	\oplus	$-(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$
Quad. lk. deg.	$1 + \langle a \rangle$	\oplus	$-(1 + \langle a \rangle)$

Or. fund. cyc.	$\eta \otimes (\overline{f_1}^* \wedge \overline{g_1}^*)$		$\eta \otimes (\overline{f_2}^* \wedge \overline{g_2}^*)$
Seifert divisors	$\langle f_1 \rangle \otimes \overline{g_1}^*$		$\langle f_2 \rangle \otimes \overline{g_2}^*$
Apply inter. prod.	$\langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x - y)$ $+ \langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x)$		
...	...		
Apply $\partial \oplus \partial$	$(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$	\oplus	$-(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$
Quad. lk. deg.	$1 + \langle a \rangle$	\oplus	$-(1 + \langle a \rangle)$

- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle b \rangle, \langle ca \rangle + \langle cb \rangle) \in W(F) \oplus W(F)$ with $a, b, c \in F^*$ such that $a + b \neq 0$. The rank modulo 2 of each component is 0.

Or. fund. cyc.	$\eta \otimes (\overline{f_1}^* \wedge \overline{g_1}^*)$		$\eta \otimes (\overline{f_2}^* \wedge \overline{g_2}^*)$
Seifert divisors	$\langle f_1 \rangle \otimes \overline{g_1}^*$		$\langle f_2 \rangle \otimes \overline{g_2}^*$
Apply inter. prod.	$\langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x - y)$ $+ \langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x)$		
...	...		
Apply $\partial \oplus \partial$	$(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$	\oplus	$-(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$
Quad. lk. deg.	$1 + \langle a \rangle$	\oplus	$-(1 + \langle a \rangle)$

- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle b \rangle, \langle ca \rangle + \langle cb \rangle) \in W(F) \oplus W(F)$ with $a, b, c \in F^*$ such that $a + b \neq 0$. The rank modulo 2 of each component is 0.
- If $F = \mathbb{R}$, the absolute value of each component is $\begin{cases} 2 & \text{if } a > 0 \\ 0 & \text{if } a < 0 \end{cases}$.

Or. fund. cyc.	$\eta \otimes (\overline{f_1}^* \wedge \overline{g_1}^*)$		$\eta \otimes (\overline{f_2}^* \wedge \overline{g_2}^*)$
Seifert divisors	$\langle f_1 \rangle \otimes \overline{g_1}^*$		$\langle f_2 \rangle \otimes \overline{g_2}^*$
Apply inter. prod.	$\langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x - y)$ $+ \langle f_1 f_2 \rangle \otimes (\overline{g_2}^* \wedge \overline{g_1}^*) \cdot (z, x)$		
...	...		
Apply $\partial \oplus \partial$	$(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$	\oplus	$-(1 + \langle a \rangle) \eta^2 \otimes (\overline{u}^* \wedge \overline{v}^*)$
Quad. lk. deg.	$1 + \langle a \rangle$	\oplus	$-(1 + \langle a \rangle)$

- If we change its orientations and its parametrizations then we get $(\langle a \rangle + \langle b \rangle, \langle ca \rangle + \langle cb \rangle) \in W(F) \oplus W(F)$ with $a, b, c \in F^*$ such that $a + b \neq 0$. The rank modulo 2 of each component is 0.
- If $F = \mathbb{R}$, the absolute value of each component is $\begin{cases} 2 & \text{if } a > 0 \\ 0 & \text{if } a < 0 \end{cases}$.
- Σ_2 of each component is $\langle a \rangle \in W(F)/(1)$. For instance, if $F = \mathbb{Q}$, Σ_2 distinguishes between all the B_p with p prime numbers. $\Sigma_4 = 0$ etc.

Everything new I presented can be found in my preprint “The quadratic linking degree”:

- HAL: Clémentine Lemarié--Rieusset. THE QUADRATIC LINKING DEGREE. 2022. ⟨hal-03821736⟩
- arXiv: Clémentine Lemarié--Rieusset. The quadratic linking degree. arXiv:2210.11048 [math.AG]

Everything new I presented can be found in my preprint “The quadratic linking degree”:

- HAL: Clémentine Lemarié--Rieusset. THE QUADRATIC LINKING DEGREE. 2022. ⟨hal-03821736⟩
- arXiv: Clémentine Lemarié--Rieusset. The quadratic linking degree. arXiv:2210.11048 [math.AG]

Thanks for your attention!